NOREDON C +
3 mg/tablet, for birds (hens broiler, turkeys)

1. NAME AND ADDRESS OF THE SALES AUTHORISATION HOLDER AND OF THE PRODUCTION AUTHORISATION HOLDER, RESPONSIBLE FOR PRODUCT SERIES ISSUE, IF THEY ARE DIFFERENT

S.C. VANELLI S.R.L.
Șos.Iași-Tg-Frumos, km10, village Letcani,
com.Letcani, jud. Iași, România 707280
T: 0040 232 253 442, F: 0040 232 253 456
E: office@vanellivet.ro, www.vanellivet.ro

2.NAME OF THE VETERINARY MEDICINAL PRODUCT :
NOREDON C+3 mg/TABLET, for birds (hens, broiler, turkeys)

3.ACTIVE SUBSTANCES AND OTHER INGREDIENTS DECLARATION :
One tablet contains: active substance: Florfenicol 3 mg

4. INDICATIONS : Birds (hens, broiler, turkeys) to treat and prevent (where the diagnosis was confirmed) bacteria infections produced by bacteria sensible to florfenicol :
-chronic respiratory disease (Mycoplasma gallisepticum)
- infectious synovitis (Mycoplasma synoviae)
- airsacculitis (Mycoplasma meleagridis)
- colibacillosis (Escherichia coli)
- avian cholera(Pasteurella multocida)
- turkey bordetella (Bordetella avium)
- infectious coryza(Haemophillus paragallinarum)

5. CONTRAINDICATIONS : Do not administer to animals sensitive to florefenicol or to other excipients. Do not use in case of known resistance to florfenicol.

6. ADVERSE REACTIONS : The most often observed reactions are diarrhea, perianal infalmmation and rectal prolapse. These symptoms are transitional and disappear after the treatment is interrupted.

If severe reactions appear or any other effects which are not mentioned in this leaflet, please inform the veterinary doctor.

7.TARGET SPECIES: Birds (hens, , broiler, turkeys)

8.POSOLOGY FOR EACH SPECIES, WAYS AND ADMINISTRATION
METHOD : Oral use.
Birds : (hens, , broiler, turkeys) : one tablet/ body weight/ day
Treatment period : 5-7 days.

9. RECOMMENDATONS FOR A CORRECT USE:
To avoid under-dosing and to ensure a correct dosage the exact body weight must be determined.

10. WAITING TIME : Birds ((hens, , broiler, turkeys) : meat and organs : 28 days. Do not administer to birds that lay eggs for human consumption.

11.SPECIAL PRECAUTIONS OF STORAGE : The product is to be kept in dry, dark place at temperatures below 25°C.

12. SPECIAL WARNINGS : Special warnings for each target species : do not exist.

Special cautions for usage to animals : When using this product keep in mind the official and local politics regarding antimicrobials. This product will be used according to isolated bacteria breakpoints. If this is not possible, the therapy will be based on epidemiologic information regarding the isolated bacteria breakpoints to animals, and also regarding the national official antimicrobial politics. Using the product outside its instructions from RCP can lead to the prevalent increase of the resistance to florfenicol and it can lead to the decrease of the efficiency of the antimicrobial treatments.

Special cautions which must be taken by the person who administers the medicinal veterinary product to animals : the persons with known hypersensitivity to florfenicol or to the excipients of the product must avoid contact. Cutaneous hypersensitivity can appear. Avoid skin contact. One must not eat, drink or smoke during the administration of the treatment. After the usage of this veterinary medicinal product , wash hands carefully with water and soap. In case of exposure , wash with plenty of water. In case of being swallowed by accident or in case

symptoms like dermal pruritus or skin irritations appear, immediately contact the doctor and show him the product leaflet and label.

Usage during gestation , lactation and in laying-eggs-period : Do not administer to birds that lay eggs for human consumption.

Interactions with other medicinal products or other forms of interactions :
No reported interactions with other medicinal products were registered to animals.

Overdosage (symptoms , urgency procedures, antidotes), according to case :
It is recommended not to exceed the therapeutical dose .

Incompatibilities : In the absence of compatibility studies , this medicinal veterinary product must not be mixed with other medicinal veterinary products.

13. Special precautions to the disposal of the un-used product or waste, according to case :
The medicine must not be thrown in residual water or household garbage.
Ask reference to the veterinary doctor regarding the way to dispose the medicine that is not necessary anymore. These measures contribute to the environmental protection.

14. INFORMATION ACCORDING TO WHICH THE LEAFLET WAS LAST TIME APPROVED : 07.2018

15. OTHER INFORMATION : Package dimensions :
Polyethylene high-density bottles x 25 tablets, x 100 tablets, x 1000 tablets.
Not all package dimensions can be sold.
For any other information regarding this veterinary medicinal product please contact the local representative who own the Sales Autorisation.